

BRASSICA

BETTER GENETICS.
BETTER PRODUCTIVITY.
BETTER PROFITABILITY.



VIVANT


Hybrid Brassica



(Vivant on left, Purple Top on right)

KEY CHARACTERISTICS

- Less Bolting
- High digestibility
- Fast establishing
- High yielding, leafy forage
- Good for grazing and attractive food plot
- Strong regrowth
- Provides quick forage
- Very leafy hybrid that has 80% more leaf than Purple Top Turnips, providing more feed for the animal



AGRONOMIC RECOMMENDATIONS

- Crops should have at least 35-55 lb phosphate/acre available at sowing.
- After grazing, a light application of nitrogen will greatly increase yields of subsequent growth.
- Avoid high nitrate levels in the crop. Paddocks should have low soil sulphate levels.

Planting Rate (direct drill)	4 lbs./acre
Planting Rate (broadcast)	6 lbs./acre
Days to Grazing	42 - 65 days
Planting Depth	1/8 to 1/4 inch deep
Potential Yield	10,000 lbs./acre
Average Yield	8,000 lbs./acre
Minimum soil pH	5.6

APPLICATION

Vivant can be broadcast seeded followed by harrowing, packing or livestock hoofing it in. Vivant may be aerial seeded in late summer for fall/winter grazing.

MANAGEMENT RECOMMENDATIONS

Allow at least 35– 40 days before initial grazing of Vivant and then expect to re-graze in 25-30 day intervals. Graze no lower than 4” if for multiple grazings. Overgrazing can damage the crown and should be avoided. To maximize regrowth potential, graze before the plant bolts.

